arXiv:1110.3887v2 [math.GT] 1 Nov 2011

$\bar{\mu}$ INVARIANT OF NANOPHRASES

YUKA KOTORII

Abstract. Two link diagrams are link homotopic if one can be transformed into the other by a sequence of Reidemeister moves and self crossing changes. Milnor introduced an invariant under link homotopy called $\bar{\mu}$. In this paper, we extend the link homotopy to nanophrases corresponding to virtual link and $\bar{\mu}$ invariant to nanophrases.

1. Introduction

A word will be a sequence of symbols, called letters, belonging to a given set $\mathcal{A}$, called alphabet. Turaev developed the theory of words based on the analogy with curves on the plane, knots in the 3-sphere, virtual knot, etc. in [8, 9, 10, 11].

A Gauss word is a sequence of letters with the condition that any letter appearing in the sequence does so exactly twice. A Gauss word can be obtained from an oriented virtual knot diagram introduced by Kauffman in [4]. Given a diagram, label the real crossings and pick a base point on the curve somewhere away from any of the real crossings. Starting from the base point, we follow the curve and read off the labels of the crossings as we pass through them. When we return to the base point, we will have a sequence of letters in which each label of a real crossing appears exactly twice. Thus this sequence is a Gauss word. It is natural to introduce combinatorial moves on Gauss words to generate equivalence relation, which we call homotopy, based on Reidemeister moves on a knot diagram.

We then assign to each crossing some data that it is under or over and so on. This leads us to the notion of nanoword by Turaev in [11]. By introducing refined combinatorial moves on nanowords, the notion of homotopy can be refined also. From this viewpoint, homotopy of Gauss words is the simplest kind of nanoword homotopy. In fact, homotopic nanowords in any refinement are homotopic as Gauss words.

The theory of nanowords can be naturally generalized to the theory of nanophrases as the knot theory does to the link theory. The purpose of this paper is to develop weaker homotopy theory on nanophrases, called M-homotopy, which is an analogue of Milnor's link homotopy [6, 7]. By a link homotopy we mean a deformotion of one link onto another, during which each component of the link is allowed to cross itself, but no two components are allowed to intersect. Milnor introduced an invariant under link homotopy called $\bar{\mu}$ in [6, 7]. We introduce a self crossing move on nanophrases and the associated M-homotopy allowing self crossings. The main result stated in Theorem 3.3 is to define an M-homotopy invariant of nanophrases corresponding to virtual links as an extension of Milnor's $\bar{\mu}$ invariant. Dye and Kauffman have done a similar work in [1]. We will come back to their results in Example 4.5. Also, Kravchenko and Polyak have done an extension of Milnor's μ invariants to virtual tangles in [5]. Their extension coincides with ours for the case

---

Date: November 2, 2011.

2000 *Mathematics Subject Classification.* Primary 57M99; Secondary 68R15.

Key words and phrases. nanowords, nanophrases, homotopy, $\bar{\mu}$ invariants.

of virtual links, modulo lower degree invariants. However our proof is considerably different from theirs.

This paper is organized as follows. In section 2, following Turaev, we give formal definitions of words, phrases and so on. In section 3, we develop M-homotopy theory for nanophrase and define invariants analogous to Milnor's $\bar{\mu}$. In section 4, we introduce some examples for $\bar{\mu}$. In section 5, 6 and 7, we prove the invariance under an M-homotopy corresponding to virtual links. In section 8, we show that this invariant is also an invariant under an M-homotopy corresponding to welded links.

2. Nanowords and nanophrases

In this section, following Turaev [8, 9, 10, 11], we review formal definitions of words, phrases and so on.

2.1. **Words and phrases.** An *alphabet* is a finite set and its element is called a *letter.* For any positive integer m, let $\hat{m}$ denote the set $\{1, 2, \dots, m\}$. A *word* on an alphabet $\mathcal{A}$ of length m is a map

$$w : \hat{m} \to \mathcal{A}.$$

Informally, we can think of a word w on $\mathcal{A}$ as a finite sequence of letters in $\mathcal{A}$ and we will usually write words in this way. For example, ABA is a word of length 3 on $\{A, B, C\}$, where 1 is mapped to A, 2 to B and 3 to A. By convention, the empty word of length 0 on any alphabet is written by $\emptyset$.

Concatenation of two words is defined by writing down the first word and then the second one. For example, the concatenation of the words $w = ABC$ and $v = DBBA$ is the word $wv = ABCDBBA$.

An n-component *phrase* on an alphabet $\mathcal{A}$ is a sequence of n words on $\mathcal{A}$. We write phrases as a sequence of words, separated by '$|$'. For example, $A|BCA|CD$ is a 3-component phrase on $\{A, B, C, D\}$. There is a unique phrase with 0-components which we denote by $\emptyset_P$. In this paper, we will regard words as 1-component phrases.

2.2. **Nanowords and nanophrases.** Let α be a finite set. An α-alphabet is an alphabet $\mathcal{A}$ together with an associated map from $\mathcal{A}$ to α. This map is called a *projection.* The image of any $A \in \mathcal{A}$ in α will be denoted by $|A|$. An *isomorphism* of α-alphabets $\mathcal{A}_1$ and $\mathcal{A}_2$ is a bijection f from $\mathcal{A}_1$ to $\mathcal{A}_2$ such that $|f(A)|$ is equal to $|A|$ for any letter A in $\mathcal{A}_1$.

A *Gauss word* on an alphabet $\mathcal{A}$ is a word on $\mathcal{A}$ such that every letter in $\mathcal{A}$ appears exactly twice. Similarly, a Gauss phrase on $\mathcal{A}$ is a phrase on $\mathcal{A}$ such that the concatenation of the words appearing in the phrase is a Gauss word on $\mathcal{A}$. By definition, a 1-component Gauss phrase is a Gauss word.

A *nanowrod* over α is a pair $(\mathcal{A}, w)$ where $\mathcal{A}$ is an α-alphabet and w is a Gauss word on $\mathcal{A}$. An n-component *nanophrase* over α is a pair $(\mathcal{A}, p)$ where $\mathcal{A}$ is an α-alphabet and p is an n-component Gauss phrase on $\mathcal{A}$.

Two nanophrases over α, $(\mathcal{A}_1, p_1)$ and $(\mathcal{A}_2, p_2)$, are *isomorphic* if there exists a bijection f from $\mathcal{A}_1$ to $\mathcal{A}_2$ such that f applied letterwise to the ith component of p_1 gives the ith component of p_2 for all i.

Rather than writing $(\mathcal{A}, p)$, we will simply use p to indicate a nanophrase. When we write a nanophrase in this way, we do not forget the set $\mathcal{A}$ of letters and the projection : $\mathcal{A} \to \alpha$.

2.3. **Equivalence relations on nanophrases.** Fix α and then let τ be an involution on α (that is, $\tau(\tau(a))$ is equal to a for all $a \in \alpha$). Let S be a subset of $\alpha \times \alpha \times \alpha$. We call the triple (α, τ, S) a *homotopy* data.

Fixing a homotopy data (α, τ, S), we define three *homotopy moves* on nanophrases over α as follows. In the moves on nanophrases, the lower cases x, y, z and t represent arbitrary sequences of letters, possibly including one or more '$|$', so that the phrase on each side of the move is a nanophrase. The moves are

move H1: for any $|A|$,

$$(\mathcal{A}, xAAy) \longleftrightarrow (\mathcal{A} - \{A\}, xy)$$

move H2: if $\tau(|A|) = |B|$,

$$(\mathcal{A}, xAByBAz) \longleftrightarrow (\mathcal{A} - \{A, B\}, xyz)$$

move H3: if $(|A|, |B|, |C|) \in S$,

$$(\mathcal{A}, xAByACzBCt) \longleftrightarrow (\mathcal{A}, xBAyCAzCBt).$$



FIGURE 1. H1



FIGURE 2. H2



FIGURE 3. H3

The *homotopy* is an equivalence relation on nanophrases over α generated by isomorphisms and three homotopy moves. The homotopy depends on the choice of the homotopy data (α, τ, S), so different choices of homotopy data give different equivalence relations. As none of the moves add or remove components, the number of components of a nanophrase is an invariant under any kind of homotopy.

In [9], Turaev defined a *shift move* on nanophrases. Let ν be an involution on α which is independent of τ. Let p be an n-component nanophrase over α. A *shift* move on the ith component of p is a move which gives a new nanophrase p' as follows. If the ith component of p is empty or contains a single letter, p' is p. If not, the ith component of p has the form Ax. Then the ith component of p' is xA and for all j not equal to i, the jth component of p' is the same as the jth component of p. Furthermore, if we write $|A|_p$ for $|A|$ in p and $|A|_{p'}$ for $|A|$ in p', then $|A|_{p'}$ equals $\nu(|A|_p)$ when x contains the letter A and otherwise, $|A|_{p'}$ equals $|A|_p$.

We call the equivalence relation generated by isomorphisms, homotopy moves and shift moves also a *homotopy* of nanophrases over α. We call the first homotopy without shift moves an *open homotopy* and the second homotopy simply a *homotopy.* The homotopy depends on the triple (α, τ, S) and ν.

Let α_v be the set $\{a_+, a_-, b_+, b_-\}$ and τ_v the involution on α_v which sends a_+ to b_- and a_- to b_+. Let S_v be the set

$$S_v = \left\{ \begin{array}{l} (a_+, a_+, a_+), (a_+, a_+, a_-), (a_+, a_-, a_-), \\ (a_-, a_-, a_-), (a_-, a_-, a_+), (a_-, a_+, a_+), \\ (b_+, b_+, b_+), (b_+, b_+, b_-), (b_+, b_-, b_-), \\ (b_-, b_-, b_-), (b_-, b_-, b_+), (b_-, b_+, b_+) \end{array} \right\}.$$

In [9], Turaev proved

Theorem 2.1 (Turaev [9]). *(i)The set of homotopy classes of nanowords over α_v under the open homotopy with respect to (α_v, τ_v, S_v) is in bijective correspondence with open virtual knots. Also under the same homotopy data, the set of homotopy classes of nanophrases over α_v is in bijective correspondence with the set of stable equivalence classes of ordered pointed link diagrams on oriented surfaces.*

(ii)Let ν_v be the involution on α_v where a_+ is mapped to b_+ and a_- to b_-. Under the homotopy defined by (α_v, τ_v, S_v) and ν_v, the set of homotopy classes

of nanophrases over α_v *is in bijective correspondence with ordered virtual links (namely, virtual links where the components are ordered and the equivalence of ordered virtual links respects the order).*

3. $\bar{\mu}$ INVARIANTS OF NANOPHRASES

In this section, we introduce a self crossing move on nanophrases and a new weak homotopy, which we call the M-homotopy, by allowing self crossing moves. Consider the set of equivalence classes of nanophrases corresponding to ordered virtual links, that is equivalence classes of nanophrases defined by (α_v, τ_v, S_v) and ν_v. We define a M-homotopy invariant of nanophrases corresponding to ordered virtual links. This invariant is an extension of $\bar{\mu}$ invariants introduced by Milnor in [6, 7].

3.1. **Self crossing and M-homotopy.** Let (α, τ, S) be any homotopy data and ν an involution on α independent of τ. We introduce the *self crossing move* on nanophrases over α. Let σ be an involution on α which is independent of τ and ν. Let p be an n-component nanophrase over α. A *self crossing move* on the kth component of p is a move which gives a new nanophrase p' as follows. If there is a letter A in $\mathcal{A}$ which appears exactly twice in the kth component of p, then the kth component of p has the form $xAyAz$. Then the kth component of p' also has the form $xAyAz$. Furthermore, writing $|A|_p$ for $|A|$ in p and $|A|_{p'}$ for $|A|$ in p', we have the identity $|A|_{p'} = \sigma(|A|_p)$.

We define the *open M-homotopy* to be the equivalence relation on nanophrases over α generated by isomorphisms, three homotopy moves with respect to (α, τ, S) and self crossing moves with respect to σ. We also define the *M-homotopy* to be the equivalence relation of nanophrases over α generated by isomorphisms, three homotopy moves with respect to (α, τ, S), self crossing moves with respect to σ and shift moves with respect to ν.

We introduce an example. The linking matrix of a nanophrase was first defined by Fukunaga [2] for some special S and by Gibson [3] in general. It is an invariant under any kind of homotopy of nanophrases. We recall the definition here. First of all, let π be the abelian group generated by elements in α with the relations $a + \tau(a) = 0$ for all a in α. In [2], π is written multiplicatively, but here we will write it additively. For an n-component nanophrase p, the *linking matrix* of p is defined as follows. Let $\mathcal{A}_{ij}(p)$ be the set of letters which have one occurrence in the ith component of p and the other occurrence in the jth component of p. Let $l_{ii}(p)$ be 0, and when j is not equal to i, let $l_{ij}(p)$ be

$$l_{ij}(p) = \sum_{A \in \mathcal{A}_{ij}(p)} |A|.$$

Let the linking matrix $L(p)$ be the symmetric $n \times n$ matrix given by $l_{ij}(p)$'s. It is easy to see that the linking matrix of nanophrases is a homotopy and M-homotopy invariant of nanophrases.

3.2. **Definition of $\bar{\mu}$.** From now on, we work only on the homotopy defined by (α_v, τ_v, S_v), ν_v and σ_v which sends a_+ to a_- and b_+ to b_-.

Let $(\mathcal{A}, p)$ be an n-component nanophrase whose Gauss phrase p is represented by $w_1|w_2|\cdots|w_n$. We will write p for $(\mathcal{A}, p)$ for simplicity. Let w_i be $A_{i1}A_{i2}\cdots A_{im_i}$, where A_{ij}'s are letters in $\mathcal{A}$. To each w_i, we define a word on $\mathcal{A} \cup \mathcal{A}^{-1}$ by

$$w_i^{\varepsilon} = A_{i1}^{\varepsilon_{i1}} A_{i2}^{\varepsilon_{i2}} \cdots A_{im_i}^{\varepsilon_{im_i}},$$

where ε_{ij} is determined as follows. Since p is a nanophrase, any letter appears exactly twice in a Gauss phrase $w_1|w_2|\cdots|w_n$. Let A denote the letter represented by A_{ij}. Then there exist integers k and l $((k,l) \neq (i,j))$ such that A_{kl} represents A. In other words, the other A appears as the lth letter on the kth component. If $i < k$ and $|A| = b_+$ or $i > k$ and $|A| = a_+$, then $\varepsilon_{ij} = 1$. If $i < k$ and $|A| = a_-$ or $i > k$ and $|A| = b_-$, then $\varepsilon_{ij} = -1$. Otherwise, $\varepsilon_{ij} = 0$. Namely, if the letter A appears exactly once in the ith component, and if A_{ij} appears earlier (or latter) than the other A and $|A| = b_+$(or a_+), then ε_{ij} is 1. If A appears exactly once in the ith component, and if A_{ij} appears earlier (or latter) than the other A and $|A| = a_-$(or b_-), then ε_{ij} is -1. For other cases, let ε_{ij} be zero. In the following, we use the convention that $A^0 = \emptyset$ and $(A_1A_2\cdots A_n)^{-1} = A_n^{-1}A_{n-1}^{-1}\cdots A_1^{-1}$. We note that $AA^{-1} \neq \emptyset$.

Let $\mathcal{L}$ denote the set of words on $A \cup A^{-1}$. Then we define a sequence of maps ρ^q $(q = 2, 3, \cdots)$ from $\mathcal{L}$ to itself by induction on q.

$$\begin{aligned}
\rho^2(A_{ij}^{\pm}) &= A_{ij}^{\pm} \\
\rho^q(A_{ij}^{\pm}) &= \rho^{q-1}(x_{ij}^{-1})A_{ij}^{\pm}\rho^{q-1}(x_{ij}),\ q \geq 3 \\
\rho^q(\emptyset) &= \emptyset \text{ for all } q \geq 2,
\end{aligned}$$

where

$$x_{ij} = A_{k1}^{\varepsilon_{k1}}A_{k2}^{\varepsilon_{k2}}\cdots A_{kl}^{\varepsilon_{kl}}.$$

Here, k and l are derived from i and j as in the above. We naturally extend ρ^q to $\mathcal{L}$. We will concern exclusively with $\rho^q(w_i^\varepsilon)$.

Example 3.1. Consider a nanophrase $p = AB|CDB|DEA|FFCE$ where $|A| = |B| = |C| = b_+, |D| = b_-, |E| = |F| = a_-$. Then

$$\begin{aligned}
&\rho^2(w_1^\varepsilon) = AB,\ \rho^2(w_2^\varepsilon) = C,\ \rho^2(w_3^\varepsilon) = D^{-1}E^{-1},\ \rho^2(w_4^\varepsilon) = \emptyset. \\
&\rho^3(w_1^\varepsilon) = EDAD^{-1}E^{-1}C^{-1}BC,\ \rho^3(w_2^\varepsilon) = C, \\
&\rho^3(w_3^\varepsilon) = C^{-1}D^{-1}CE^{-1},\ \rho^3(w_4^\varepsilon) = \emptyset. \\
&\rho^4(w_1^\varepsilon) = EC^{-1}DCAC^{-1}D^{-1}CE^{-1}C^{-1}BC,\ \rho^4(w_2^\varepsilon) = C, \\
&\rho^4(w_3^\varepsilon) = C^{-1}D^{-1}CE^{-1},\ \rho^4(w_4^\varepsilon) = \emptyset. \\
&\cdots\cdots\cdots
\end{aligned}$$

Recall that $p = w_1|w_2|\cdots|w_n$ is an n-component nanophrase. We call the index i of a component w_i the order of w_i. Let M denote a finite set $\{a_1, \ldots, a_n\}$. Let $\mathcal{M}$ denote the set of words on $M \cup M^{-1}$. Using a nanophrase property of p, we define a map η from $\mathcal{L}$ to $\mathcal{M}$ as follows. For any letter A in $\mathcal{A}$, let $\eta(A)$ be a_k and $\eta(A^{-1})$ a_k^{-1}, where k is determined by the following rule. If $|A| = b+$ or $a-$, then k is the order of the component in p in which the second A occurs, and if $|A| = a+$ or $b-$, then k is the order of the component in p in which the first A occurs. To see what k is for a letter in w_i, let us recall the definition of w_i^ε. The letter A in w_i survives in w_i^ε only when either $|A| = b_+$ or a_- and the other A appears in latter component, or $|A| = a_+$ or b_- and the other appears in former component. Thus, if A is represented by A_{ij} and $\varepsilon_{ij} \neq 0$, then k is the number appeared in the definition of ε_{ij}, namely, the order of the component in which the other A occurs.

Example 3.2. Consider a nanophrase $p = AB|CDB|DEA|FFCE$ where $|A| = |B| = |C| = b_+$, $|D| = b_-$, $|E| = |F| = a_-$ (example 3.1). Then we have

$$\eta(\rho^2(w_1^\varepsilon)) = a_3a_2,\ \eta(\rho^2(w_2^\varepsilon)) = a_4,\ \eta(\rho^2(w_3^\varepsilon)) = a_2^{-1}a_4^{-1},\ \eta(\rho^2(w_4^\varepsilon)) = \emptyset.$$
$$\eta(\rho^3(w_1^\varepsilon)) = a_4a_2a_3a_2^{-1}a_4^{-1}a_4^{-1}a_2a_4,\ \eta(\rho^3(w_2^\varepsilon)) = a_4,$$
$$\eta(\rho^3(w_3^\varepsilon)) = a_4^{-1}a_2^{-1}a_4a_4^{-1},\ \eta(\rho^3(w_4^\varepsilon)) = \emptyset.$$
$$\eta(\rho^4(w_1^\varepsilon)) = a_4a_4^{-1}a_2a_4a_3a_4^{-1}a_2^{-1}a_4a_4^{-1}a_4^{-1}a_2a_4,\ \eta(\rho^4(w_2^\varepsilon)) = a_4,$$
$$\eta(\rho^4(w_3^\varepsilon)) = a_4^{-1}a_2^{-1}a_4a_4^{-1},\ \eta(\rho^4(w_4^\varepsilon)) = \emptyset.$$
$$\cdots\cdots\cdots$$

We define a map φ from $\mathcal{M}$ to $\mathbb{Z}[[\kappa_1, \kappa_2, \dots, \kappa_n]]$ by

$$\begin{aligned}\varphi(a_h) &= 1 + \kappa_h,\\ \varphi(a_h^{-1}) &= 1 - \kappa_h + \kappa_h^2 - \kappa_h^3 + \cdots,\end{aligned}$$

where $\mathbb{Z}[[\kappa_1, \kappa_2, \dots, \kappa_n]]$ is the ring of formal power series on non-commuting variables $\kappa_1, \kappa_2, \dots, \kappa_n$.

We consider $\varphi \circ \eta(\rho^q(w_i^\varepsilon))$ in $\mathbb{Z}[[\kappa_1, \kappa_2, \dots, \kappa_n]]$. Since $\varphi \circ \eta(\rho^q(w_i^\varepsilon))$ agrees with $\varphi \circ \eta(\rho^r(w_i^\varepsilon))$ for any $r \geq q$ up to degree q, the coefficient of a term $\kappa_{c_1}\kappa_{c_2}\dots\kappa_{c_u}$ in $\varphi \circ \eta(\rho^q(w_i^\varepsilon))$ converges as $q \to \infty$. Thus we have a well-defined expansion,

$$\lim_{q\to\infty} \varphi \circ \eta(\rho^q(w_i^\varepsilon)) = 1 + \sum \mu(p; c_1, c_2, \dots, c_u, i)\kappa_{c_1}\kappa_{c_2}\cdots\kappa_{c_u}.$$

where $c_1, c_2, \dots, c_u, i$ is a sequence of integers between 1 and n. We here note that the integers in the sequence are not necessarily mutually different.

Let $\Delta(p; c_1, c_2, \dots, c_u, i)$ denote the greatest common divisor of $\mu(p; d_1, d_2, \dots, d_t)$, where the sequence $d_1, d_2, \dots, d_t$ $(2 \leq t \leq u)$ ranges over all sequences obtained by eliminating at least one of $c_1, c_2, \dots, c_u, i$, and permuting the remaining indices cyclically. We also define $\Delta(p; c_1, i) = 0$. Let $\bar{\mu}(p; c_1, c_2, \dots, c_u, i)$ denote the residue class of $\mu(p; c_1, c_2, \dots, c_u, i)$ modulo $\Delta(p; c_1, c_2, \dots, c_u, i)$.

The main theorem of this paper is as follows.

Theorem 3.3. *Let p be an n-component nanophrase. Let $c_1, c_2, \dots, c_u, i$ be a sequence of integers between* 1 *and n such that $c_1, c_2, \dots, c_u, i$ are pairwise distinct. Then $\bar{\mu}(p; c_1, c_2, \dots, c_u, i)$ is an invariant under M-homotopy of nanophrases with respect to (α_v, τ_v, S_v), ν_v and σ_v.*

Proposition 6.1, 6.2, 6.3, 6.4 and 7.4 will imply the proof.

4. Example

Example 4.1. Let $p = ABCD|ECFA|DFBE$ where $|A| = |E| = b+$, $|B| = b-$, $|C| = |F| = a-$ and $|D| = a+$. This corresponds to the Borromean rings illustrated in Fig. 4. Then since $w_1^\varepsilon = AC^{-1}$,

$$\begin{aligned}&\rho^2(w_1^\varepsilon) = AC^{-1}\\ &\rho^3(w_1^\varepsilon) = FE^{-1}AEF^{-1}E^{-1}C^{-1}E.\end{aligned}$$

Thus we have

$$\begin{aligned}&\eta(\rho^2(w_1^\varepsilon)) = a_2a_2^{-1}\\ &\eta(\rho^3(w_1^\varepsilon)) = a_3a_3^{-1}a_2a_3a_3^{-1}a_3^{-1}a_2^{-1}a_3.\end{aligned}$$

Therefore $\mu(2,1) = \mu(3,1) = 0$ and $\mu(2,3,1) = -1$. Similarly $\mu(1,2) = \mu(1,3) = \mu(2,3) = \mu(3,2) = 0$ and so $\Delta(2,3,1) = 0$. Hence $\bar{\mu}(2,3,1) \equiv -1 \pmod 0$.



FIGURE 4.

Example 4.2. Let $p = A_1B_1A_2B_2 \dots A_nB_n|A_1B_1A_2B_2 \dots A_nB_n$ where $|A_j| = b+$ and $|B_j| = a+$. This corresponds to the $(2n, 2)$ torus link illustrated in Fig. 5. Then since $w_1^\varepsilon = A_1 \dots A_n$ and $w_2^\varepsilon = B_1 \dots B_n$,

$$\eta(\rho^2(w_1^\varepsilon)) = a_2 \dots a_2$$
$$\eta(\rho^2(w_2^\varepsilon)) = a_1 \dots a_1.$$

Therefore $\mu(2,1) = n$ and $\mu(1,2) = n$, and so $\bar{\mu}(2,1) = n$ and $\bar{\mu}(1,2) = n$.



FIGURE 5.

Example 4.3. Let $p = A_1A_2 \dots A_n|A_1A_2 \dots A_n$ where $|A_j| = b+$. This corresponds to the virtual link illustrated in Fig. 6. Then since $w_1^\varepsilon = A_1 \dots A_n$ and $w_2^\varepsilon = \emptyset$,

$$\eta(\rho^2(w_1^\varepsilon)) = a_2 \dots a_2$$
$$\eta(\rho^2(w_2^\varepsilon)) = \emptyset.$$

Therefore $\mu(2,1) = n$ and $\mu(1,2) = 0$, and so $\bar{\mu}(2,1) = n$ and $\bar{\mu}(1,2) = 0$.



FIGURE 6.

Example 4.4. Let $p = AB|FBCDAE|DC|EF$ where $|A| = |B| = |E| = b+$, $|C| = b-$, $|D| = a-$ and $|F| = a+$. This corresponds to the link illustrated in Fig. 7. Then since $w_1^\varepsilon = AB$,

$$\rho^2(w_1^\varepsilon) = AB$$
$$\rho^3(w_1^\varepsilon) = DAD^{-1}B.$$

Thus we have

$$\eta(\rho^2(w_1^\varepsilon)) = a_2a_2$$
$$\eta(\rho^3(w_1^\varepsilon)) = a_3a_2a_3^{-1}a_2.$$

Therefore $\mu(2,1) = 2$, $\mu(3,1) = 0$, $\mu(2,3,1) = -1$ and $\mu(3,2,1) = 1$. Similarly $w_2^\varepsilon = D^{-1}E$ and $w_3^\varepsilon = C^{-1}$. Thus we have

$$\eta(\rho^2(w_2^\varepsilon)) = a_3^{-1}a_4$$
$$\eta(\rho^2(w_3^\varepsilon)) = a_2^{-1}.$$

Therefore $\mu(1,2) = 0$, $\mu(3,2) = -1$, $\mu(1,3) = 0$ and $\mu(2,3) = -1$. So we have $\Delta(2,3,1) = \Delta(3,2,1) = 1$. As a result, $\bar{\mu}(2,3,1) \equiv \bar{\mu}(3,2,1) \equiv 0 \pmod 1$.



Figure 7.

Example 4.5. Let $p = ABCD|DAEFBC|EF|\emptyset$ where $|A| = |C| = |F| = a+$, $|B| = |D| = |E| = b+$. This corresponds to the link illustrated in Fig. 8. Then since $w_1^\varepsilon = BD$,

$$\rho^2(w_1^\varepsilon) = BD$$
$$\rho^3(w_1^\varepsilon) = E^{-1}A^{-1}BAED.$$

Thus we have

$$\eta(\rho^2(w_1^\varepsilon)) = a_2a_2$$
$$\eta(\rho^3(w_1^\varepsilon)) = a_3^{-1}a_1^{-1}a_2a_1a_3a_2.$$

Therefore $\mu(2,1) = 2$, $\mu(3,1) = 0$ and $\mu(2,3,1) = 1$. Similarly $w_2^\varepsilon = AEC$ and $w_3^\varepsilon = F$. Thus we have

$$\eta(\rho^2(w_2^\varepsilon)) = a_1a_3a_1$$
$$\eta(\rho^2(w_3^\varepsilon)) = a_2.$$

Therefore $\mu(1,2) = 2$, $\mu(3,2) = 1$, $\mu(1,3) = 0$ and $\mu(2,3) = 1$. So we have $\Delta(2,3,1) = 1$. As a result, $\bar{\mu}(2,3,1) \equiv 0 \pmod 1$.



Figure 8.

Example 4.5 made us wonder if there is something missing in the proof of homotopy invariance of $\bar{\mu}$-invariant in Dye and Kauffman [1].

In fact, Example 4.5 exhibits an example of a classical link where their $\bar{\mu}$ depends on the choice of base points. If we choose a base point so that p in Example 4.5 is the corresponding phrase, then their $\mu(p;2,1) = 2$, $\mu(p;3,1) = 0$ and $\mu(p;2,3,1) = 1$, and therefore their $\bar{\mu}(p;2,3,1) \equiv 1 \bmod 2$. On the other hand, if we replace the base point of the second component so that the corresponding new phrase is $p' = ABCD|FBCDAE|EF|\emptyset$ as in Fig. 9, then their $\mu(p';2,1) = 2$, $\mu(p';3,1) = 0$ and $\mu(p';2,3,1) = 0$, and hence we have their $\bar{\mu}(p';2,3,1) \equiv 0 \bmod 2$.



Figure 9.

Example 4.5 exhibits an example of a classical link where the indeterminacy Δ depends on the choice of the base point. If we choose a base point so that p in Example 4.5 is the corresponding phrase, then their $\mu(p;2,3,1) = 1$ and in particular $\Delta(p;2,3,4,1) = 1$. On the other hand, if we replace the base point of the second component so that the corresponding new phrase is $p' = ABCD|FBCDAE|EF|\emptyset$, then their $\mu(p';2,1) = 2$, $\mu(p';3,1) = \mu(p';4,1) = \mu(p';2,3,1) = \mu(p';3,2,1) = \mu(p';2,4,1) = \mu(p';4,2,1) = \mu(p';3,4,1) = \mu(p';4,3,1) = 0$, and hence we have their $\Delta(p';2,3,4,1) = 2$.

We should comment also that Dye and Kauffman's $\bar{\mu}$ differs from Milnor's in some classical case. Let L be the link illustrated in Fig. 10. By Dye and Kauffman's definition, $\mu(1,3) = \mu(2,3) = 0$ and hence $\Delta(1,2,3) = 0$, and therefore their $\bar{\mu}(1,2,3) \equiv 0 \bmod 0$. On the other hand, Milnor's $\Delta(1,2,3) = 1$ for L, and therefore his $\bar{\mu}(1,2,3) \equiv 0 \bmod 1$.



Figure 10.

5. Preliminaries to the proof of invariance

It is sufficient to prove that for any two nanophrases p and p' related by either isomorphisms, H1 moves, H2 moves, H3 moves, shift moves or self crossing moves, for each sequence $c_1, c_2, \dots, c_u, i$ of distinct integers between 1 and n, $\bar{\mu}(p;c_1,c_2,\dots,c_u,i)$ is equal to $\bar{\mu}(p';c_1,c_2,\dots,c_u,i)$. It is easy to see that this holds for an isomorphism. To show the invariance under the other moves we prepare graphs.

Given a nanophrase $p = w_1|w_2|\cdots|w_n$, we defined its signed ith component $w_i^{\varepsilon} = A_{i1}^{\varepsilon_{i1}} A_{i2}^{\varepsilon_{i2}} \cdots A_{im_i}^{\varepsilon_{im_i}}$ in the previous section.

We will construct a sequence of rooted trees $T_{ij}^q = T_{ij}^q(p) = T_{A_{ij}}^q(p)$ for each $A_{ij}^{\varepsilon_{ij}}$ and a sequence of forests $F_i^q = F_i^q(p)$ by assembling T_{ij}^q's, where q is the integer associated with the expanding word $\rho^q(A_{ij}^{\varepsilon_{ij}})$ of $A_{ij}^{\varepsilon_{ij}}$. To do this, recall how we defined $\rho^q(A_{ij}^{\varepsilon_{ij}})$ from $\rho^{q-1}(A_{ij}^{\varepsilon_{ij}})$. Since the operation of ρ^q on $\rho^{q-1}(A_{ij}^{\varepsilon_{ij}})$ is to insert several words in $\rho^{q-1}(A_{ij}^{\varepsilon_{ij}})$, $\rho^{q-1}(A_{ij}^{\varepsilon_{ij}})$ is obtained from $\rho^q(A_{ij}^{\varepsilon_{ij}})$ by deleting some subwords. If we denote this inclusive relation by $<$, we have a natural inclusion of words,

$$\rho^2(A_{ij}^{\varepsilon_{ij}}) < \rho^3(A_{ij}^{\varepsilon_{ij}}) < \cdots < \rho^q(A_{ij}^{\varepsilon_{ij}})$$

Then, we assign two natural numbers to each letter in the word $\rho^q(A_{ij}^{\varepsilon_{ij}})$. One is the depth. To each letter in $\rho^q(A_{ij}^{\varepsilon_{ij}})$, we assign the depth d if it appears in $\rho^{d+2}(A_{ij}^{\varepsilon_{ij}})$ but not in $\rho^{d+1}(A_{ij}^{\varepsilon_{ij}})$. The depth is an intrinsic invariant for a letter in $\rho^q(A_{ij}^{\varepsilon_{ij}})$ independent from q. The other is its location in $\rho^q(A_{ij}^{\varepsilon_{ij}})$. We assign an integer g to the gth letter in $\rho^q(A_{ij}^{\varepsilon_{ij}})$. Note that the location g depends on q. Also g distinguishes vertices in T_{ij}^q. We now define an increasing sequence of rooted trees,

$$T_{ij}^2 \subset T_{ij}^3 \subset \cdots \subset T_{ij}^q \subset \cdots \tag{1}$$

The vertices of T_{ij}^q consists of letters in $\rho^q(A_{ij}^{\varepsilon_{ij}})$. To each vertex, we assign its depth. We join two vertices v and w by a directed edge from v to w if the depth d of v is equal to the depth of w minus 1 and moreover w is a letter appeared in $\rho^3(v)$ but not in $\rho^2(v)$. Thus we get a sequence of rooted trees (1) based on the letter $A_{ij}^{\varepsilon_{ij}}$ in w_i^{ε}. The root of each tree is represented uniquely by $A_{ij}^{\varepsilon_{ij}}$ of depth 0.

For further discussion, we assign a label to each vertices in T_{ij}^q. It is a quadruple consisting of the location g of the letter in $\rho^q(A_{ij}^{\varepsilon_{ij}})$ where the letter in question appears as the gth letter, its letter in $\mathcal{A}$ without a sign ε, the index k of its image under η, and the sign ε of its letter. If we regard T_{ij}^q as a labeled rooted tree, then, the vertices can be distinguished by the first factor. On the other hand, the inclusive relation in (1) still dose make sense if we ignore the first factor of the labels.

We now denote the forest by labeled rooted trees $T_{ij}^q(p)$'s by $F_i^q(p)$, where j ranges between 1 and m_i. Note that $F_i^q(p)$ contains m_i roots of depth 0.

Example 5.1. Consider the nanophrase $p = AB|CDB|DEA|FFCE$ where $|A| = |B| = |C| = b_+, |D| = b_-, |E| = |F| = a_-$ (example 3.1). Then we have

$$\rho^2(w_1^\varepsilon) = AB,\ \rho^3(w_1^\varepsilon) = EDAD^{-1}E^{-1}C^{-1}BC,$$
$$\rho^4(w_1^\varepsilon) = EC^{-1}DCAC^{-1}D^{-1}CE^{-1}C^{-1}BC,$$
$$\eta(\rho^4(w_1^\varepsilon)) = a_4a_4^{-1}a_2a_4a_3a_4^{-1}a_2^{-1}a_4a_4^{-1}a_4^{-1}a_2a_4.$$

Then $F_1^4(p)$ is the following forest.



Let F be a forest by rooted trees with labels in $\mathbb{N} \times \mathcal{A} \times \{1, 2, \cdots, n\} \times \{\pm 1\}$ such that vertices of each forest can be distinguished by the first factor of the label. Then, for any sequence $c_1, c_2, \cdots, c_u$ of pairwise different integers between 1 and n, let $\mathcal{S}(F, c_1, \cdots, c_u)$ be the set of subforests of F satisfying the following conditions:

(1) Each member of $\mathcal{S}(F, c_1, c_2, \cdots, c_u)$ has u vertices.
(2) If their first factors are labeled by $d_1, d_2, \cdots, d_u$ such that $d_1 < d_2 < \cdots < d_u$, then the third factor of the vertex labeled by d_j is c_j.
(3) Each component contains a root.

We then define sets $\mathcal{S}_e(F, c_1, \cdots, c_u)$ and $\mathcal{S}_o(F, c_1, \cdots, c_u)$ as follows. For any sequence $c_1, c_2, \cdots, c_u$ of pairwise different integers between 1 and n, let $\mathcal{S}_e(F, c_1, \cdots, c_u)$

and $\mathcal{S}_o(F, c_1, \cdots, c_u)$ be subsets of $\mathcal{S}(F, c_1, \cdots, c_u)$ which consists of subforests for which the cardinality of the vertices with $\varepsilon_j = -1$ is even or odd respectively. Note that ε_j is the sign of the forth factor. Also note that we have the identity

$$\mathcal{S}(F, c_1, \cdots, c_u) = \mathcal{S}_o(F, c_1, \cdots, c_u) \cup \mathcal{S}_e(F, c_1, \cdots, c_u).$$

Example 5.2. Consider the nanophrase $AB|CDB|DEA|FFCE$ where $|A| = |B| = |C| = b_+, |D| = b_-, |E| = |F| = a_-$ (example 3.1). Then we have the following where we note that we write only the first factor of the label.

$\mathcal{S}(L_1^4, 3, 2, 4) =$



$\mathcal{S}_e(L_1^4, 3, 2, 4) =$



$\mathcal{S}_o(L_1^4, 3, 2, 4) =$



Lemma 5.3. *If the indices $c_1, c_2, \ldots, c_u$ are pairwise distinct, then for any i and q the coefficient of $\kappa_{c_1} \cdots \kappa_{c_u}$ in $\varphi \circ \eta(\rho^q(w_i^\varepsilon))$ is equal to the cardinality of $\mathcal{S}_e(F_i^q, c_1, \ldots, c_u)$ minus the cardinality of $\mathcal{S}_o(F_i^q, c_1, \ldots, c_u)$.*

Proof. We consider the coefficient of $\kappa_{c_1} \cdots \kappa_{c_u}$ in $\varphi \circ \eta(\rho^q(w_i^\varepsilon))$. $\varphi \circ \eta(\rho^q(w_i^\varepsilon))$ is the product of some elements of the form $(1 + \kappa_c)$ and $(1 - \kappa_d + {\kappa_d}^2 - {\kappa_d}^3 + \cdots)$ in $\mathbb{Z}[[\kappa_1, \kappa_2, \ldots, \kappa_n]]$, where $1 \leq c, d \leq n$. Let E denote the set of all power series $\sum \lambda(e_1, e_2, \ldots, e_v)\kappa_{e_1}\kappa_{e_2} \ldots \kappa_{e_v}$ in $\mathbb{Z}[[\kappa_1, \kappa_2, \ldots, \kappa_n]]$ such that at least two indices in $\{e_1, e_2, \ldots e_v\}$ coincide. It is easy to see that E is a two-sided ideal. Since the indices $c_1, c_2, \ldots, c_u$ are pairwise distinct by assumption, it is sufficient to consider the coefficient of $\kappa_{c_1} \cdots \kappa_{c_u}$ modulo E. Therefore we regard $1 - \kappa + \kappa^2 - \kappa^3 + \cdots$ as $1 - \kappa$ modulo E. On the other hand, $\rho^q(w_i^\varepsilon)$ is the product of $\rho^q(A_{ij}^{\varepsilon_{ij}})$ $(1 \leq j \leq m_i)$. Moreover by the definition of ρ,

$$\rho^q(A_{ij}^{\varepsilon_{ij}}) = \rho^{q-1}(x_{ij}^{-1})A_{ij}^{\varepsilon_{ij}}\rho^{q-1}(x_{ij}).$$

Then the letter with depth 0 in $\rho^q(A_{ij}^{\varepsilon_{ij}})$ is only $A_{ij}^{\varepsilon_{ij}}$ $(1 \leq j \leq m_i)$ in the right hand of this identity. Let $\eta(A_{ij}) = a_{e_j}$,

$$\varphi \circ \eta(A_{ij}^{\varepsilon_{ij}}) \equiv 1 + \varepsilon_{ij}\kappa_{e_j} \pmod{E}.$$

In addition, let y_j denote $\varphi \circ \eta(\rho^{q-1}(x_{ij}))$. Then

(2)
$$\varphi \circ \eta(\rho^q(w_i^\varepsilon)) \equiv y_1^{-1}(1 + \varepsilon_{i1}\kappa_{e_1})y_1 \cdot y_2^{-1}(1 + \varepsilon_{i2}\kappa_{e_2})y_2 \cdots y_{m_i}^{-1}(1 + \varepsilon_{im_i}\kappa_{e_{m_i}})y_{m_i} \pmod{E}.$$

First of all, we expand all the factor $(1 + \varepsilon_{ij}\kappa_{e_j})$ in (2). Then since the number of letters with depth 0 is m_i, (2) is expanded into the 2^{m_i} terms. Secondly, to each terms we cancel out $y_j^{-1}1y_j$ for any j. We repeat these two steps according to depths. Then it follows from how to expand the expression and how expansion constructs the forest F_i^q that there exists the natural bijection between terms in this expansion and subforests of F_i^q, where each component contains a root. The coefficient of each term in this expanded expression is ± 1. On the other hand, for each term in this expansion, we consider the vertices of the subforest corresponding to it. The number of vertices with -1 as the forth factor of the labels modulo 2 corresponds to the coefficient of the term in this expansion. If the number of this vertices is even (odd), then the coefficient is 1 (-1, respectively). Therefore

$$\begin{aligned}\text{the coefficient of } &\kappa_{c_1}\cdots\kappa_{c_u} \text{ in } \varphi\circ\eta(\rho^q(w_i^\varepsilon))\\ &= \#\mathcal{S}_{\mathrm{e}}(F_i^q, c_1, \dots, c_u) - \#\mathcal{S}_{\mathrm{o}}(F_i^q, c_1, \dots, c_u).\end{aligned}$$

□

Example 5.4. Consider a nanophrase $p = AB|CDB|DEA|FFCE$ where $|A| = |B| = |C| = b_+, |D| = |E| = b_-, |F| = a_-$ (example 3.1). Then we have

$$\begin{aligned}\eta(\rho^3(w_3^\varepsilon)) &= a_4^{-1}a_2^{-1}a_4a_4^{-1},\\ \varphi\circ\eta(\rho^3(w_3^\varepsilon)) &= (1-\kappa_4+\kappa_4^2-\cdots)(1-\kappa_2+\kappa_2^2-\cdots)(1+\kappa_4)(1-\kappa_4+\kappa_4^2-\cdots).\end{aligned}$$

Therefore

$$\varphi\circ\eta(\rho^3(w_3^\varepsilon))$$

$$\equiv (1-\kappa_4)(1-\kappa_2)(1+\kappa_4)(1-\kappa_4) \pmod{E} \tag{3}$$

$$\begin{aligned}&= (1-\kappa_4)1(1+\kappa_4)1 + (1-\kappa_4)1(1+\kappa_4)(-\kappa_4)\\ &\qquad + (1-\kappa_4)(-\kappa_2)(1+\kappa_4)1 + (1-\kappa_4)(-\kappa_2)(1+\kappa_4)(-\kappa_4)\end{aligned} \tag{4}$$

$$\equiv 1-\kappa_4 + (1-\kappa_4)(-\kappa_2)(1+\kappa_4) + (1-\kappa_4)(-\kappa_2)(1+\kappa_4)(-\kappa_4) \pmod{E} \tag{5}$$

$$\begin{aligned}&= 1-\kappa_4 + \Big(-\kappa_2 - \kappa_2\kappa_4 + \kappa_4\kappa_2 + \kappa_4\kappa_2\kappa_4\Big)\\ &\qquad\qquad + \Big(\kappa_2\kappa_4 + \kappa_2\kappa_4\kappa_4 - \kappa_4\kappa_2\kappa_4 - \kappa_4\kappa_2\kappa_4\kappa_4\Big).\end{aligned} \tag{6}$$

where for (3), expanding the terms corresponding to letters with depth 0, we obtain (4). Moreover for (4), simplifying the expansion by canceling out them we obtain (5). In addition to (5), expanding the terms corresponding to letters with depth 1, we obtain (6). We note that the coefficient of terms in (6) is ±1.

Here we have

$F_3^3(p) =$



On the other hand, (6) corresponds to the subforests as follows where we note that we write only the first factor of the label.



Then, for example, the sum of the coefficient of $\kappa_2\kappa_4$ in (6) is 0. On the other hand, since $\#\mathcal{S}_{\mathrm{e}}(F_i^q, c_1, \dots, c_u)$ and $\#\mathcal{S}_{\mathrm{o}}(F_i^q, c_1, \dots, c_u)$ are 1,

$$\#\mathcal{S}_{\mathrm{e}}(F_i^q, c_1, \dots, c_u) - \#\mathcal{S}_{\mathrm{o}}(F_i^q, c_1, \dots, c_u) = 0$$

6. Proof of invariance for $H1, H2, H3$ and self crossing moves

Proposition 6.1. *If the indices $c_1, c_2, \ldots, c_u, i$ are a sequence of integers between 1 and n, then $\mu(p; c_1, \ldots, c_u, i) = \mu(p'; c_1, \ldots, c_u, i)$ if p and p' are related by H1 move.*

Proof. Let p' be $w'_1|w'_2|\cdots|w'_n$. Suppose $p = xAAy$ and $p' = xy$. The letter A appears twice in a certain component. It follows from the definition of $\rho^q(w_i^\varepsilon)$ that the letter A is not contained in $\rho^q(w_i^\varepsilon)$ and thus does not affect $\rho^q(w_i^\varepsilon)$. Thus $\rho^q(w_i^\varepsilon)$ is equal to $\rho^q({w'_i}^\varepsilon)$, and so for any sequence $c_1, c_2, \ldots, c_u, i$, $\mu(p; c_1, c_2, \ldots, c_u, i)$ is equal to $\mu(p'; c_1, c_2, \ldots, c_u, i)$. ☐

Proposition 6.2. *If the indices $c_1, c_2, \ldots, c_u, i$ are a sequence of integers between 1 and n, then $\mu(p; c_1, \ldots, c_u, i) = \mu(p'; c_1, \ldots, c_u, i)$ if p and p' are related by H2 move.*

Proof. Suppose $p = xAByBAz$, where $|A|$ is equal to $\tau_v(|B|)$ and $p' = xyz$. We need to consider the following two cases.

Case 1: The subwords AB and BA appear in a certain component.
(i.e. p is a nanophrase of the form $\ldots|\ldots AB \ldots BA \ldots|\ldots$)
Case 2: The subwords AB and BA appear in different components.
(i.e. p is a nanophrase of the form $\ldots|\ldots AB \ldots|\ldots|\ldots BA \ldots|\ldots$)

Case1: The letter A and B appear twice in a certain component. It follows from the definition of $\rho^q(w_i^\varepsilon)$ that the letters A and B are not contained in $\rho^q(w_i^\varepsilon)$ and thus do not affect $\rho^q(w_i^\varepsilon)$. Thus $\rho^q(w_i^\varepsilon)$ is equal to $\rho^q({w'_i}^\varepsilon)$, and $\mu(p; c_1, \ldots, c_u, i)$ is equal to $\mu(p'; c_1, \ldots, c_u, i)$.

Case2: Let h be the order of the component in which the subword AB occurs, k the order of the component in which the subword BA occurs. We show that $\varphi \circ \eta(\rho^q(w_i^\varepsilon))$ is equal to $\varphi \circ \eta(\rho^q({w'_i}^\varepsilon))$. We suppose that $|A| = b_+$, then $|B| = a_-$. (We can show the others cases in the same way.) Then the sign of A in hth component is 1 and the sign of B in hth component -1. Moreover, the signs of A and B in kth component are 0, and they do not appear directly in $\rho^q(w_i^\varepsilon)$. Since A and B in hth component are adjacent and their signs are ± 1, either $\rho^q({w'_i}^\varepsilon) = \rho^q(w_i^\varepsilon)$ when $\rho^r(w_i^\varepsilon)$ contains neither subwords AB^{-1} nor BA^{-1} for any r $(2 \leq r \leq q)$, or $\rho^q({w'_i}^\varepsilon)$ is obtained from $\rho^q(w_i^\varepsilon)$ by deleting subwords $\rho^r(AB^{-1})$ and $\rho^r(BA^{-1})$ for some $2 \leq r \leq q$. Since A and B in kth component are adjacent and their signs are 0,

$$\begin{aligned}\rho^r(AB^{-1}) &= \rho^r(A)\rho^r(B^{-1}) \\ &= \rho^{r-1}(B^0x^{-1})A\rho^{r-1}(xB^0)\rho^{r-1}(x^{-1})B^{-1}\rho^{r-1}(x) \\ &= \rho^{r-1}(x^{-1})A\rho^{r-1}(x)\rho^{r-1}(x^{-1})B^{-1}\rho^{r-1}(x).\end{aligned}$$

By $\eta(A) = a_k$ and $\eta(B^{-1}) = a_k^{-1}$,

$$\begin{aligned}\eta(\rho^r(AB^{-1})) &= \eta(\rho^{r-1}(x^{-1})A\rho^{r-1}(x)\rho^{r-1}(x^{-1})B^{-1}\rho^{r-1}(x)) \\ &= \eta(\rho^{r-1}(x^{-1}))a_k\eta(\rho^{r-1}(x))\eta(\rho^{r-1}(x^{-1}))a_k^{-1}\eta(\rho^{r-1}(x)).\end{aligned}$$

Therefore

$$\varphi \circ \eta(\rho^r(AB^{-1})) = 1.$$

Similarly $\varphi \circ \eta(\rho^r(BA^{-1})) = 1$. Therefore $\varphi \circ \eta(\rho^q(w_i^\varepsilon))$ is equal to $\varphi \circ \eta(\rho^q({w'_i}^\varepsilon))$. Thus $\mu(p; c_1, \ldots, c_u, i)$ is equal to $\mu(p'; c_1, \ldots, c_u, i)$. ☐

Proposition 6.3. *If the indices $c_1, c_2, \ldots, c_u, i$ are pairwise distinct integers between 1 and n, then $\mu(p; c_1, \ldots, c_u, i) = \mu(p'; c_1, \ldots, c_u, i)$ if p and p' are related by H3 move.*

Proof. Suppose $p = xAByACzBCt$, where $(|A|, |B|, |C|)$ is an element of S and $p' = xBAyCAzBCt$. Moreover we suppose that $|A| = |B| = |C| = b_+$. (We can show the others cases in the same way.) We need to consider the following four cases.

Case 1: The subwords AB, AC and BC appear in a certain component.
(i.e. p is a nanophrase of the form $...|...AB...AC...BC...|...$)

Case 2: The subwords AB and AC appear in a certain component, but the subword BC does not appear in that component.
(i.e. p is a nanophrase of the form $...|...AB...AC...|...|...BC...|...$)

Case 3: The subwords AC and BC appear in a certain component, but the subword AB does not appear in that component.
(i.e. p is a nanophrase of the form $...|...AB...|...|...AC...BC...|...$)

Case 4: The subwords AB, AC and BC appear in three different components.
(i.e. p is a nanophrase of the form $...|...AB...|...|...AC...|...|...BC...|...$)

Case1: The letters A, B and C appear twice in a component and hence it does not affect $\rho^q(w_i^\varepsilon)$ and $\rho^q({w'_i}^{\varepsilon})$. Thus $\rho^q(w_i^\varepsilon)$ is equal to $\rho^q({w'_i}^{\varepsilon})$.

Case2: The letter A appears twice in a component and hence it does not affect $\rho^q(w_i^\varepsilon)$. Since in the component which contains the subword BC the letters B and C are adjacent and the sign associated with these letters are 0, they do not appear directly in $\rho^q(w_i^\varepsilon)$. Thus $\rho^q(w_i^\varepsilon)$ is equal to $\rho^q({w'_i}^{\varepsilon})$.

Case3: Let h be the order of the component in which the subword AB occurs, k the order of the component in which the subwords AC and BC occur. By the same reason as in Case 1, the letter C does not affect $\rho^q(w_i^\varepsilon)$ and $\rho^q({w'_i}^{\varepsilon})$. We divide the proof into the following two cases.

Case 3-1: $i = k$
Case 3-2: $i \neq k$

Case 3-1: Consider the subgraph F (F', respectively) of $F_k^q(p)$ ($F_k^q(p')$, respectively) induced by the set of vertices obtained by deleting the vertices labeled by k and their descendants. If we ignore the first factor of the labels, then the subgraphs F and F' are isomorphic as forests by labeled rooted trees. Since $i = k$, $k \notin \{c_1, ..., c_u\}$ by assumption. So the cardinality of $\mathcal{S}_{\mathrm{e}}(F_k^q(p), c_1, \dots, c_u)$ is equal to the cardinality of $\mathcal{S}_{\mathrm{e}}(F, c_1, \dots, c_u)$, and the same identity holds for $F_k^q(p')$ and F' as well. It follows from lemma 5.3 that

$$\begin{aligned}
\mu(p; c_1, \dots, c_u, k) &= \#\mathcal{S}_{\mathrm{e}}(F_k^q(p), c_1, \dots, c_u) - \#\mathcal{S}_{\mathrm{o}}(F_k^q(p), c_1, \dots, c_u) \\
&= \#\mathcal{S}_{\mathrm{e}}(F, c_1, \dots, c_u) - \#\mathcal{S}_{\mathrm{o}}(F, c_1, \dots, c_u) \\
&= \#\mathcal{S}_{\mathrm{e}}(F', c_1, \dots, c_u) - \#\mathcal{S}_{\mathrm{o}}(F', c_1, \dots, c_u) \\
&= \#\mathcal{S}_{\mathrm{e}}(F_k^q(p'), c_1, \dots, c_u) - \#\mathcal{S}_{\mathrm{o}}(F_k^q(p'), c_1, \dots, c_u) \\
&= \mu(p'; c_1, \dots, c_u, k).
\end{aligned}$$

Case 3-2: If $k \notin \{c_1, \dots, c_u\}$, it follows from the same reason as in case 3-1 that $\mu(p) = \mu(p')$. Let $k \in \{c_1, \dots, c_u\}$. We divide $\mathcal{S}(F_i^q(p), c_1, \dots, c_u)$ into three groups according to whether it contains a vertex labeled by A or B or not. Let $\mathcal{G}_{e,A}(p)$ denote the set of elements of $\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \dots, c_u)$ whose vertices do not have A as the second factor of the labels. Let $\mathcal{G}_{e,B}(p)$ denote the set of elements of $\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \dots, c_u)$ whose vertices do not have B as the second factor of the labels. Let $\mathcal{G}_e(p)$ denote the set of elements of $\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \dots, c_u)$ whose vertices labeled by neither A nor B. Similarly, we define $\mathcal{G}_{o,A}(p)$, $\mathcal{G}_{o,B}(p)$ and $\mathcal{G}_o(p)$. Since $\eta(A) = \eta(B) = a_k$, we have

$$\begin{aligned}
\mu(p; c_1, \dots, c_u, i) &= \#\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \dots, c_u) - \#\mathcal{S}_{\mathrm{o}}(F_i^q(p), c_1, \dots, c_u) \\
&= \#\mathcal{G}_{e,A}(p) - \#\mathcal{G}_{o,A}(p) + \#\mathcal{G}_{e,B}(p) - \#\mathcal{G}_{o,B}(p) - (\#\mathcal{G}_e(p) - \#\mathcal{G}_o(p)).
\end{aligned}$$

The above identity also holds if we replace p by p'. Consider the subgraph F_A (F'_A, respectively) of $F_i^q(p)$ ($F_i^q(p')$, respectively) induced the set of vertices obtained by deleting every vertices labeled by A and their descendants. Similarly, consider the subgraph F_B (F'_B, respectively) of $F_i^q(p)$ ($F_i^q(p')$, respectively) induced by the set of vertices obtained by deleting every vertices labeled by B and their descendants. Similarly, consider the subgraph F (F', respectively) of $F_i^q(p)$ ($F_i^q(p')$, respectively) induced by the set of vertices obtained by deleting every vertices labeled by either A or B and their descendants. Then we have

$$\mathcal{G}_{e,A}(p) = \mathcal{S}_{\mathrm{e}}(F_A, c_1, \dots, c_u)$$

and so on. If we ignore the first factor of the labels, then F_A and F'_A are isomorphic as forests by labeled rooted trees. Moreover F_B and F'_B, and F and F' are also isomorphic. Therefore we have

$$\begin{aligned}\#\mathcal{S}_{\mathrm{e}}(F_A, c_1, \dots, c_u) &= \#\mathcal{S}_{\mathrm{e}}(F'_A, c_1, \dots, c_u),\\ \#\mathcal{S}_{\mathrm{e}}(F_B, c_1, \dots, c_u) &= \#\mathcal{S}_{\mathrm{e}}(F'_B, c_1, \dots, c_u),\\ \#\mathcal{S}_{\mathrm{e}}(F, c_1, \dots, c_u) &= \#\mathcal{S}_{\mathrm{e}}(F', c_1, \dots, c_u),\end{aligned}$$

and so on. Thus $\mu(p; c_1, \dots, c_u, i)$ is equal to $\mu(p'; c_1, \dots, c_u, i)$.

Case 4: Let h be the order of the component in which the subword AB occurs, j the order of the component in which the subword AC occurs, and k the order of the component in which the subword BC occurs. We divide the proof into the following three cases.

Case 4-1: $i = j$
Case 4-2: $i = k$
Case 4-3: $i \neq j, k$

Case 4-1: Consider the subgraph of $F_j^q(p)$ ($F_j^q(p')$) induced by the set of virtices obtained by deleting vertices labeled by A and their descendants. If we ignore the first factor of the labels, then they are isomorphic as forests by labeled rooted trees. Since $j \notin \{c_1, \dots, c_u\}$ and $\eta(A) = a_j$, it follows from the case 3-1 that we have $\mu(p; c_1, \dots, c_u, j) = \mu(p'; c_1, \dots, c_u, j)$.

Case 4-2: Consider the subgraph of $F_i^q(p)$ ($F_i^q(p')$, respectively) induced by the set of virtices obtained by deleting vertices labeled by B and C and their descendants. If we ignore the first factor of the label, then they are isomorphic as forests by labeled rooted trees. Since $k \notin \{c_1, \dots, c_u\}$ and $\eta(B) = \eta(C) = a_k$, we have $\mu(p; c_1, \dots, c_u, k) = \mu(p'; c_1, \dots, c_u, k)$.

Case 4-3: We divide $\mathcal{S}(F_i^q(p), c_1, \dots, c_u)$ into two groups according to whether it contains a vertex with A as a second factor of the labels or not. Let $\mathcal{H}_{e,A}(p)$ denote the set of elements of $\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \dots, c_u)$ whose vertices contain a vertex labeled by A. Let $\mathcal{G}_{e,A}(p)$ denote as before the set of elements of $\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \dots, c_u)$ whose vertices do not contain a vertex labeled by A. Similarly, we define $\mathcal{H}_{o,A}$ and $\mathcal{G}_{o,A}$. Then we have

$$\begin{aligned}\mu(p; c_1, \dots, c_u, i) &= \#\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \dots, c_u) - \#\mathcal{S}_{\mathrm{o}}(F_i^q(p), c_1, \dots, c_u)\\ &= \#\mathcal{H}_{e,A}(p) - \#\mathcal{H}_{o,A}(p) + \#\mathcal{G}_{e,A}(p) - \#\mathcal{G}_{o,A}(p).\end{aligned} \tag{7}$$

Consider the right-hand side of the equality in (7). We denote the sum of first two terms by (7.1) and the last two terms by (7.2). First of all, we consider (7.2). Consider the subgraph of $F_i^q(p)$ ($F_i^q(p')$, respectively) induced by the set of vertices obtained by deleting vertices labeled by A and their descendants. If we ignore the first factor of the label, then they are isomorphic as forests by labeled rooted trees. Therefore we have

$$(7.2) = \#\mathcal{G}_{e,A}(p') - \#\mathcal{G}_{o,A}(p'). \tag{8}$$

Secondly, we consider (7.1). Let V_A denote the set of all vertices of a forest $F_i^q(p)$ labeled by A which does not have ancestor labeled by A. Fix an element v in V_A. Then we consider the set of vertices obtained from vertices of $F_i^q(p)$ by deleting the following vertices. If the depth of v is $q-2$, the vertices are all vertices of $F_i^q(p)$ labeled by A except v, their descendants and the vertex labeled by B paired with v, where it means that the vertex which has the same parent as v, has B as the second factor of the labels and has the same forth factor of the labels as v. If not, the vertices are all vertices of $F_i^q(p)$ labeled by A except v, their descendants and all vertices of depth $q-2$ which are ancestors of the vertex labeled by B paired with v. Let F_v denote the subgraph of $F_i^q(p)$ induced by them. Similarly V'_A denote the set of all vertices of a forest $F_i^q(p')$ labeled by A which does not have ancestor labeled by A. There exists a natural bijection from V_A to V'_A. Fix the element v' in V'_A to which the bijection maps v. Then let $F'_{v'}$ denote the subgraph of $F_i^q(p')$ as above. If the depth of v is $q-2$, then it is easy to see that the subword of $\rho^q(w_i^\varepsilon)$ and $\rho^q({w'_i}^\varepsilon)$ corresponding to F_v and F'_v, respectively are equal. If the depth of v is $s<q-2$, then v, the vertex labeled by B paired with v and their descendants in F_v correspond to the subword $\rho^{q-s}(A)\rho^{q-s-1}(B)$ or $\rho^{q-s-1}(B^{-1})\rho^{q-s}(A^{-1})$. Then let

$$\rho^{q-s}(A)\rho^{q-s-1}(B)=\rho^{q-s-1}(x^{-1})A\rho^{q-s-1}(x)\rho^{q-s-1}(B). \tag{9}$$

Moreover v', the vertex labeled by B paired with v' and their descendants in $F'_{v'}$ correspond to the subword $\rho^{q-s-1}(B)\rho^{q-s}(A)$ or $\rho^{q-s}(A^{-1})\rho^{q-s-1}(B^{-1})$. Then we have

$$\rho^{q-s-1}(B)\rho^{q-s}(A)=\rho^{q-s-1}(B)\rho^{q-s-1}(C^{-1})\rho^{q-s-1}(x^{-1})A\rho^{q-s-1}(x)\rho^{q-s-1}(C). \tag{10}$$

Since $\eta(\rho^r(B))=\eta(\rho^r(C))$ for any r, the image of (9) and (10) under $\varphi\circ\eta$ are equal. Similarly, the image of $\rho^{q-s-1}(B^{-1})\rho^{q-s}(A^{-1})$ and $\rho^{q-s}(A^{-1})\rho^{q-s-1}(B^{-1})$ under $\varphi\circ\eta$ are equal. Therefore the image of the subword of $\rho^q(w_i^\varepsilon)$ corresponding to F_v under $\varphi\circ\eta$ is equal to the image of the subword of $\rho^q({w'_i}^\varepsilon)$ corresponding to $F'_{v'}$ under $\varphi\circ\eta$. Thus we have

$$\begin{aligned}(7.1)&=\sum_{v\in V_A}\Big(\#\{G\in\mathcal{S}_{\mathrm{e}}(F_i^q(p),c_1,\dots,c_u)\mid v\in V(G)\}\\&\qquad\qquad-\#\{G\in\mathcal{S}_{\mathrm{o}}(F_i^q(p),c_1,\dots,c_u)\mid v\in V(G)\}\Big)\\&=\sum_{v\in V_A}\Big(\#\mathcal{S}_{\mathrm{e}}(F_v,c_1,\dots,c_u)-\#\mathcal{S}_{\mathrm{o}}(F_v,c_1,\dots,c_u)\Big)\\&=\sum_{v'\in V'_A}\Big(\#\mathcal{S}_{\mathrm{e}}(F'_{v'},c_1,\dots,c_u)-\#\mathcal{S}_{\mathrm{o}}(F'_{v'},c_1,\dots,c_u)\Big)\\&=\sum_{v'\in V'_A}\Big(\#\{G\in\mathcal{S}_{\mathrm{e}}(F_i^q(p'),c_1,\dots,c_u)\mid v'\in V(G)\}\\&\qquad\qquad-\#\{G\in\mathcal{S}_{\mathrm{o}}(F_i^q(p'),c_1,\dots,c_u)\mid v'\in V(G)\}\Big)\end{aligned} \tag{11}$$

where $V(G)$ is the set of vertices of G. Therefore we have

$$(8)+(11)=\mu(p';c_1,\dots,c_u,i).$$

$\square$

Proposition 6.4. *If the indices $c_1,c_2,\dots,c_u,i$ are a sequence of integers between 1 and n, then $\mu(p;c_1,\dots,c_u,i)=\mu(p';c_1,\dots,c_u,i)$ if p and p' are related by self crossing move.*

Proof. Suppose $p = ...|...A...A...|...$, and p' also has the same form. Furthermore, if we write $|A|_p$ for $|A|$ in p and $|A|_{p'}$ for $|A|$ in p', $|A|_{p'}$ equals $\sigma_v(|A|_p)$. Since A appear twice in a component, $\rho^q(w_i^\varepsilon)$ is equal to $\rho^q(w_i'^\varepsilon)$. □

7. Proof of invariance for shift move

We will show that $\mu(p; c_1, c_2, \ldots, c_u, i)$ is invariant under the shift move by induction on u. We suppose that if for each $u < q-1$ the indices $c_1, c_2, \ldots, c_u, i$ are pairwise distinct, then $\bar{\mu}(p; c_1, c_2, \ldots, c_u, i)$ is invariant under M-homotopy. Moreover we suppose that phrases p and p' are M-homotopic, then $\bar{\mu}(p; c_1, c_2, \ldots, c_u, i)$ is congruent to $\bar{\mu}(p'; c_1, c_2, \ldots, c_u, i)$ for $u < q-1$. Therefore it follows from the definition of Δ that $\Delta(p; c_1, c_2, \ldots, c_u, i)$ is congruent to $\Delta(p'; c_1, c_2, \ldots, c_u, i)$ for $u < q$. Let $D_i^q = D_i^q(p)$ denote the set of all power series $\sum \nu(c_1, c_2, \ldots, c_u)\kappa_{c_1}\kappa_{c_2}\cdots\kappa_{c_u}$ in $\mathbb{Z}[[\kappa_1, \kappa_2, \ldots, \kappa_n]]$ whose coefficients satisfy

$$\nu(c_1, c_2, \ldots, c_u) \equiv 0 \pmod{\Delta(p; c_1, c_2, \ldots, c_u, i)}$$

for all $c_1, c_2, \ldots, c_u$ with $u < q$. (There is no restriction on $\nu(c_1, c_2, \ldots, c_u)$ for $u \geq q$.)

Proposition 7.1 (Milnor [7]). *D_i^q is a two-sided ideal.*

Proof. In fact, if $\nu(c_1, c_2, \ldots, c_u)\kappa_{c_1}\kappa_{c_2}\cdots\kappa_{c_u}$ is a monomial in D_i^q, and $\lambda\kappa_{d_1}\kappa_{d_2}\cdots\kappa_{d_t}$ is an arbitrary monomial, then either $u \geq q$ so that the product is trivial in D_i^q, or

$$\nu(c_1, c_2, \ldots, c_u) \equiv 0 \pmod{\Delta(p; c_1, c_2, \ldots, c_u, i)}$$

and

$$\Delta(p; c_1, c_2, \ldots, c_u, i) \equiv 0 \pmod{\Delta(p; d_1, \ldots d_t, c_1, \ldots, c_u, i)},$$

hence

$$\lambda\nu(c_1, c_2, \ldots, c_u) \equiv 0 \pmod{\Delta(p; d_1, \ldots d_t, c_1, \ldots, c_u, i)};$$

which implies that

$$\lambda\nu(c_1, c_2, \ldots, c_u)\kappa_{d_1}\ldots\kappa_{d_t}\kappa_{c_1}\ldots\kappa_{c_u} \in D_i^q.$$

This proves that D_i^q is a left ideal. The fact that D_i^q is a right ideal is followed similarly. □

By the definition of D_i^q, $D_i^u(p)$ is congruent to $D_i^u(p')$ for any $u \leq q$. By using these hypothesises, we will show that $\bar{\mu}(p; c_1, \ldots, c_{q-1}, i)$ is equal to $\bar{\mu}(p'; c_1, \ldots, c_{q-1}, i)$.

We prepare the following two lemmas for the next proposition.

Lemma 7.2. *The image of $\rho^q(w_i^\varepsilon)$ under $\varphi \circ \eta$ is equal to the image of its any conjugate under $\varphi \circ \eta$ modulo D_i^q.*

Proof. Let $\varphi \circ \eta(\rho^q(w_i^\varepsilon)) = 1 + \omega_i$. Then $\omega_i\kappa_j \equiv \kappa_j\omega_i \equiv 0 \pmod{D_i^q}$ for any κ_j. This is proved as follows. Let $\mu(p; c_1, c_2, \ldots, c_u, i)\kappa_{c_1}\kappa_{c_2}\cdots\kappa_{c_u}$ be any term of ω_i. Then the congruence

$$\mu(p; c_1, c_2, \ldots, c_u, i) \equiv 0 \pmod{\Delta(p; c_1, c_2, \ldots, c_u, j, i)}$$

implies that

$$\mu(p; c_1, c_2, \ldots, c_u, i)\kappa_{c_1}\kappa_{c_2}\cdots\kappa_{c_u}\kappa_j \equiv 0 \pmod{D_i^q}.$$

The second congruence follows by the same method. Moreover

$$\begin{aligned}
\varphi(a_j\eta(\rho^q(w_i^\varepsilon))a_j^{-1}) &= (1+\kappa_j)(1+\omega_i)(1-\kappa_j+{\kappa_j}^2-{\kappa_j}^3+\cdots) \\
&= 1+\omega_i+(\text{ terms involving } \kappa_j\omega_i \text{ or } \omega_i\kappa_j) \\
&\equiv \varphi(\eta(\rho^q(w_i^\varepsilon))) \pmod{D_i^q}
\end{aligned}$$

where the last step follows because D_i^q is a two-side ideal. Since any conjugation can be broken down into a sequence of conjugations by a_j and a_j^{-1}, the proof is complete. □

Lemma 7.3. *For any element* a_j^ε $(\varepsilon = \pm 1)$ *in* $M \cup M^{-1}$*, we consider its conjugation by image of* $\rho^s(w_j^\varepsilon)$ *under* η *where* $s < q$*. This image under* φ *is equal to the image of* a_j^ε *under* φ *modulo* D_i^{s+1}*. That is, for any* $u \leq s < q$ *the coefficients of* $\kappa_{c_1}\kappa_{c_2}\cdots\kappa_{c_u}$ *in these two images are equal modulo* $\Delta(p; c_1, c_2, \ldots, c_u, i)$*.*

Proof. Let $\varphi \circ \eta(\rho^s(w_j^\varepsilon)) = 1 + \omega_j$. Then $\omega_j\kappa_j \equiv \kappa_j\omega_j \equiv 0 \pmod{D_i^{s+1}}$. In fact, let $\mu(p; c_1, c_2, \ldots, c_u, j)\kappa_{c_1}\kappa_{c_2}\cdots\kappa_{c_u}$ be any term of ω_j which has degree smaller than s. Then the congruences

$$\mu(p; c_1, c_2, \ldots, c_u, j) \equiv 0 \pmod{\Delta(p; c_1, c_2, \ldots, c_u, j, i)}$$

and

$$\mu(p; c_1, c_2, \ldots, c_u, j) \equiv 0 \pmod{\Delta(p; j, c_1, c_2, \ldots, c_u, i)}$$

show that

$$\mu(p; c_1, c_2, \ldots, c_u, j)\kappa_{c_1}\kappa_{c_2}\cdots\kappa_{c_u}\kappa_j \equiv 0 \pmod{D_i^{s+1}}$$

and

$$\mu(p; c_1, c_2, \ldots, c_u, j)\kappa_j\kappa_{c_1}\kappa_{c_2}\cdots\kappa_{c_u} \equiv 0 \pmod{D_i^{s+1}}.$$

For any term in ω_j which has degree greater than or equal to s, in $\omega_j\kappa_j$ and $\kappa_j\omega_j$ the degree of the product of the term and κ_j is greater than s. Thus these are zero modulo D_i^{s+1}. Moreover,

$$\begin{aligned}\varphi(a_j\eta(\rho^s(w_j^\varepsilon))) - \varphi(\eta(\rho^s(w_j^\varepsilon))a_j) &= (1+\kappa_j)(1+\omega_j) - (1+\omega_j)(1+\kappa_j)\\ &= \kappa_j\omega_j - \omega_j\kappa_j\end{aligned}$$

shows that

$$\begin{aligned}&\varphi(a_j^{-1}\eta(\rho^s(w_j^\varepsilon))^{-1}a_j\eta(\rho^s(w_j^\varepsilon)))\\ &\quad= 1 + \varphi(a_j^{-1}\eta(\rho^s(w_j^\varepsilon))^{-1})\{\varphi(a_j\eta(\rho^s(w_j^\varepsilon))) - \varphi(\eta(\rho^s(w_j^\varepsilon))a_j)\}\\ &\quad= 1 + \varphi(a_j^{-1}\eta(\rho^s(w_j^\varepsilon))^{-1})(\kappa_j\omega_j - \omega_j\kappa_j)\\ &\quad\equiv 1 \pmod{D_i^{s+1}}.\end{aligned}$$

Hence

$$\varphi(\eta(\rho^s(w_j^\varepsilon))^{-1}a_j\eta(\rho^s(w_j^\varepsilon))) = \varphi(a_j) \pmod{D_i^{s+1}}.$$

Similarly,

$$\varphi(\eta(\rho^s(w_j^\varepsilon))^{-1}a_j^{-1}\eta(\rho^s(w_j^\varepsilon))) = \varphi(a_j^{-1}) \pmod{D_i^{s+1}}.$$

□

Proposition 7.4. *If the indices* $c_1, c_2, \ldots, c_u, i$ *are pairwise distinct, then* $\bar{\mu}(p; c_1, \ldots, c_u, i) = \bar{\mu}(p'; c_1, \ldots, c_u, i)$ *if* p *and* p' *are related by the shift move.*

Proof. Suppose p and p' are related by the shift move on the kth component. That is $p = w_1|\cdots|w_{k-1}|Ax|w_{k+1}|\cdots|w_n$ and $p' = w_1|\cdots|w_{k-1}|xA|w_{k+1}|\cdots|w_n$, where A is a letter in $\mathcal{A}$. Furthermore, let we write $|A|_p$ for $|A|$ in p and $|A|_{p'}$ for $|A|$ in p', if x contains the letter A, then $|A|_{p'}$ equals $\nu_v(|A|_p)$. Otherwise, $|A|_{p'}$ equals $|A|_p$. We need to consider the following three cases.

Case 1: x contains the letter A
Case 2: $w_1\cdots w_{k-1}$ contains the letter A and $|A| = b+$ or $a-$
  (or $w_{k+1}\cdots w_n$ contains the letter A and $|A| = a+$ or $b-$)
Case 3: $w_1\cdots w_{k-1}$ contains the letter A and $|A| = a+$ or $b-$
  (or $w_{k+1}\cdots w_n$ contains the letter A and $|A| = b+$ or $a-$)

Case 1: Similar to H1 move, $\rho^q(w_i^\varepsilon)$ is equal to $\rho^q({w'}_i^{\varepsilon})$ and so $\mu(p, c_1, ..., c_{q-1}, i)$ is equal to $\mu(p', c_1, ..., c_{q-1}, i)$. Thus $\bar{\mu}(p, c_1, ..., c_{q-1}, i) = \bar{\mu}(p', c_1, ..., c_{q-1}, i)$.

Case 2: Let $|A| = b+$. (We can show the others cases in the same way.) Let k be the order of the component in which shifted A occurs and h the order of the component in which the other A occurs. We divide the proof into the following two cases.

Case 2-1: $i = k$
Case 2-2: $i \neq k$

Case 2-1: Consider the subgraph of $F_k^q(p')$ induced by the vertices obtained by deleting descendants of vertices labeled by A, and denote it by F'_A. If we ignore the first factor of the label, then $F_k^q(p)$ and F'_A are isomorphic as forests. Since $i = k$, $k \notin \{c_1, ...c_{q-1}\}$. Moreover $\eta(A) = a_k$. So the cardinality of $\mathcal{S}_{\mathrm{e}}(F_k^q(p'), c_1, \ldots, c_{q-1})$ is equal to the cardinality of $\mathcal{S}_{\mathrm{e}}(F'_A, c_1, \ldots, c_{q-1})$. Similar to the case 3-1 of $H2$ move, $\mu(p'; c_1, \ldots, c_{q-1}, k) = \mu(p; c_1, \ldots, c_{q-1}, k)$.

Case 2-2: Consider the subgraph of $F_i^q(p')$ induced by the vertices obtained by deleting descendants of vertices labeled by A. If we ignore the first factor of the label, then $F_i^q(p)$ and this subgraph are isomorphic as forests. We divide $\mathcal{S}(F_i^q(p'), c_1, \ldots, c_{q-1})$ into two groups according to whether it contains a vertex labeled by A or not. Let $\mathcal{H}_+(p')$ denote the set which consists of elements of $\mathcal{S}_{\mathrm{e}}(F_i^q(p'), c_1, \ldots, c_{q-1})$ which have a child of a vertex labeled by A. Let $\bar{\mathcal{H}}_+(p')$ denote the complement of $\mathcal{H}_+(p')$ with respect to $\mathcal{S}_{\mathrm{e}}(F_i^q(p'), c_1, \ldots, c_{q-1})$. Similarly, we define $\mathcal{H}_-(p')$ and $\bar{\mathcal{H}}_-(p')$.

$$\begin{aligned} &\mu(p'; c_1, \ldots, c_{q-1}, i) \\ &\quad = \#\mathcal{S}_{\mathrm{e}}(F_i^q(p'), c_1, \ldots, c_{q-1}) - \#\mathcal{S}_{\mathrm{o}}(F_i^q(p'), c_1, \ldots, c_{q-1}) \\ &\quad = \#\mathcal{H}_+(p') - \#\mathcal{H}_-(p') + \#\bar{\mathcal{H}}_+(p') - \#\bar{\mathcal{H}}_-(p'). \end{aligned} \tag{12}$$

Consider the right-hand side of the second equality. We denote the sum of first two terms by (12.1) and the last two terms by (12.2). First of all, we consider (12.2). $\bar{\mathcal{H}}_+(p')$ corresponds to $\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \ldots, c_{q-1})$ one-to-one and so dose $\bar{\mathcal{H}}_-(p')$ to $\mathcal{S}_{\mathrm{o}}(F_i^q(p), c_1, \ldots, c_{q-1})$. Therefore we have

$$\begin{aligned} (12.2) &= \#\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \ldots, c_{q-1}) - \#\mathcal{S}_{\mathrm{o}}(F_i^q(p), c_1, \ldots, c_{q-1}) \\ &= \mu(p; c_1, \ldots, c_{q-1}, i). \end{aligned}$$

Therefore the rest is to show that (12.1) is equal to zero modulo $\Delta(p'; c_1, \ldots, c_{q-1}, i)$. For any element G in $\mathcal{H}_+$ or $\mathcal{H}_-$, let $\hat{G}$ denote the subgraph induced by the vertices obtained from vertices of G by deleting the descendants of the vertex labeled by A. Then (12.1) is divided into the following terms.

$$(12.1) = \sum_{F \in \{\hat{G} | G \in \mathcal{H}_+ \cup \mathcal{H}_-\}} \Big(\#\{G \in \mathcal{H}_+ \mid \hat{G} = F\} - \#\{G \in \mathcal{H}_- \mid \hat{G} = F\} \Big). \tag{13}$$

Suppose that F is fixed. Let r denote the depth of the vertex of F labeled by A. Then we choose an element G in $\mathcal{H}_+$ or $\mathcal{H}_-$ such that $\hat{G}$ is F. Let a sequence $b_1, b_2, ..., b_t$ denote the subsequence of $c_1, \ldots, c_{q-1}$ obtained from $c_1, \ldots, c_{q-1}$ by picking up the sequence corresponding to the vertex labeled by A and its descendants in G. Let $s = q - r - 1$, then the right hand term of (13) is equal to

$$\sum_{F \in \{\hat{G} | G \in \mathcal{H}_+ \cup \mathcal{H}_-\}} \varepsilon_F \cdot \Big(\#\mathcal{S}_{\mathrm{e}}(T', b_1, \ldots, b_t) - \#\mathcal{S}_{\mathrm{o}}(T', b_1, \ldots, b_t) \Big) \tag{14}$$

where T' is either $T_A^{s+1}(p')$ or the tree obtained by changing the sign of the forth factor of the root in $T_A^{s+1}(p')$. We then consider the vertices of F except the vertex labeled by A. If the number of theses vertices which have label -1 as the forth factor of the labels is even (odd, respectively), let $\varepsilon_F = 1$ ($\varepsilon_F = -1$, respectively). By lemma 5.3, the right hand term of (14) is equal to

$$\sum_{F \in \{\hat{G} | G \in \mathcal{H}_+ \cup \mathcal{H}_-\}} \varepsilon_F \cdot \Big(\text{the coefficient of } \kappa_{b_1} \cdots \kappa_{b_t} \text{ in } \varphi\big(\eta(\rho^s((w_k^\varepsilon)^{-1}))a^{\pm}\eta(\rho^s(w_k^\varepsilon))\big)\Big). \tag{15}$$

We know that $r + t - 1 \leq q - 2$ and so $t \leq s$. By lemma 7.3, the coefficient of $\kappa_{b_1} \cdots \kappa_{b_t}$ in $\varphi\big(\eta(\rho^s((w_k^\varepsilon)^{-1}))a^{\pm}\eta(\rho^s(w_k^\varepsilon))\big)$ is equal to the coefficient of $\kappa_{b_1} \cdots \kappa_{b_t}$ in $\varphi(a_k^{\pm})$ modulo $\Delta(p'; b_1, b_2, \ldots, b_t, i)$, and so $\Delta(p'; c_1, \ldots, c_{q-1}, i)$. Since $t \geq 2$ and $b_1, \ldots, b_t$ are pairwise distinct, the coefficient of $\kappa_{b_1} \cdots \kappa_{b_t}$ in $\varphi(a_k^{\pm})$ is zero. Therefore term (15) is equal to zero modulo $\Delta(p'; c_1, \ldots, c_{q-1}, i)$. Thus $\bar{\mu}(p'; c_1, \ldots, c_{q-1}, i) = \bar{\mu}(p; c_1, \ldots, c_{q-1}, i)$.

Case 3: Let $|A| = a+$. (We can show the others cases in the same way.) Let k be the order of the component in which shifted A occurs, h the order of the component in which the other A. We divide the proof into the following three cases.

Case 3-1: $i = h$
Case 3-2: $i = k$
Case 3-3: $i \neq h, k$

Case 3-1: Similar to the case 4-1 of $H3$ move, $\mu(p'; c_1, \ldots, c_{q-1}, h) = \mu(p; c_1, \ldots, c_{q-1}, h)$.

Case 3-2: We can represent $\rho^2(w_k^\varepsilon)$ as Ax_2 where x_2 is a word in $\mathcal{A} \cup \mathcal{A}^{-1}$. Then $\rho^2({w'_k}^\varepsilon)$ is represented by $x_2 A$. Moreover $\rho^q(w_k^\varepsilon)$ and $\rho^q({w'_k}^\varepsilon)$ can be represented by $y_q^{-1} A y_q x_q$ and $x'_q {y'_q}^{-1} A y'_q$ respectively where x_q, y_q, x'_q and y'_q are some words in $\mathcal{A} \cup \mathcal{A}^{-1}$. Consider the set of vertices V and V' which are obtained from vertices of $F_k^q(p)$ and $F_k^q(p')$ respectively by deleting vertices labeled by k and their descendants . Let F and F' denote the two subgraphs induced by V and V' respectively. Moreover let l and l' denote the words corresponding to F and F' respectively. We note that $l < \rho^q(w_k^\varepsilon)$ and $l' < \rho^q({w'_k}^\varepsilon)$. Then l and l' can be represented by ${y''_q}^{-1} A y''_q x''_q$ and $x''_q {y''_q}^{-1} A y''_q$ respectively, where x''_q and y''_q are some words in $\mathcal{A} \cup \mathcal{A}^{-1}$. Then we have

$$\begin{aligned} \mu(p; c_1, \ldots, c_{q-1}, k) &= \#\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \ldots, c_{q-1}) - \#\mathcal{S}_{\mathrm{o}}(F_i^q(p), c_1, \ldots, c_{q-1}) \\ &= \#\mathcal{S}_{\mathrm{e}}(F, c_1, \ldots, c_{q-1}) - \#\mathcal{S}_{\mathrm{o}}(F, c_1, \ldots, c_{q-1}) \\ &= \text{the coefficient of } \kappa_{c_1} \cdots \kappa_{c_{q-1}} \text{ in } \varphi \circ \eta(l). \qquad (16) \end{aligned}$$

Similarly, we have

$$\mu(p'; c_1, \ldots, c_{q-1}, k) = \text{the coefficient of } \kappa_{c_1} \cdots \kappa_{c_{q-1}} \text{ in } \varphi \circ \eta(l'). \tag{17}$$

We notice that l is equal to $({y''_q}^{-1} A y''_q) l' ({y''_q}^{-1} A^{-1} y''_q)$. It follows from lemma 7.2 that the coefficient of $\kappa_{c_1} \cdots \kappa_{c_{q-1}}$ in $\varphi \circ \eta(l)$ is equal to that in $\varphi \circ \eta(l')$ modulo $\Delta(p'; c_1, \ldots, c_{q-1}, k)$. As a result, we have

$$\bar{\mu}(p; c_1, \ldots, c_{q-1}, k) = \bar{\mu}(p'; c_1, \ldots, c_{q-1}, k).$$

Case 3-3: $F_i^q(p')$ is the subgraph induced by the vertices obtained from vertices of $F_i^q(p)$ by deleting vertices labeled by A and thier descendants. We divide $\mathcal{S}(F_i^q(p), c_1, \ldots, c_{q-1})$ into two groups according to whether it contains a vertex of labeled by A or not. Let $\bar{\mathcal{G}}_{e,A}(p)$ denote the complement of $\mathcal{G}_{e,A}(p)$ with respect to $\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \ldots, c_{q-1})$. Similarly, we define $\bar{\mathcal{G}}_{o,A}(p)$. Then we have

$$\begin{aligned} &\mu(p; c_1, \ldots, c_{q-1}, i) \\ &\quad = \#\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \ldots, c_{q-1}) - \#\mathcal{S}_{\mathrm{o}}(F_i^q(p), c_1, \ldots, c_{q-1}) \\ &\quad = \#\bar{\mathcal{G}}_{e,A}(p) - \#\bar{\mathcal{G}}_{o,A}(p) + \#\mathcal{G}_{e,A}(p) - \#\mathcal{G}_{o,A}(p). \qquad (18) \end{aligned}$$

Consider the right-hand side of the second identity in (18). We denote the sum of first two terms by (18.1) and last two terms by (18.2). First of all, we consider (18.2). We have as before

$$(18.2) = \mu(p'; c_1, \dots, c_{q-1}, i).$$

Secondly, we consider (18.1). Since $i \neq k$, the vertices labeled by A have a vertex labeled by k as their parents. Let $\mathcal{K}_+(p)$ denote the set which consists of all elements of $\mathcal{S}_{\mathrm{e}}(F_i^q(p), c_1, \dots, c_{q-1})$ such that there exists an edge from a vertex labeled by k to a vertex labeled by A. Similarly, we define $\mathcal{K}_-(p)$. Hence

$$(18.1) = \#\mathcal{K}_+(p) - \#\mathcal{K}_-(p). \tag{19}$$

We consider the set of all sequences obtained from $c_1, \dots, c_{q-1}$ by deleting some indices where we must delete indices h and not delete indices k, and denote it by $\mathcal{C}$. For any element G of $\mathcal{S}(F_i^q(p), c_1, \dots, c_{q-1})$, $\hat{G}$ denotes the subgraph induced by the vertices obtained from the vertices of G by deleting the vertices labeled by A and their descendants. Then (19) is equal to

$$\sum_{(b_1,\dots,b_r)\in\mathcal{C}} \Big(\#\{G \in \mathcal{K}_+ \mid \hat{G} \in \mathcal{S}(F_i^q(p), b_1, ..., b_r)\} - \#\{G \in \mathcal{K}_- \mid \hat{G} \in \mathcal{S}(F_i^q(p), b_1, ..., b_r)\}\Big). \tag{20}$$

Because of the fact proved later, the term (20) is equal to

$$\sum_{(b_1,\dots b_r)\in\mathcal{C}} \delta_{b_1,\dots,b_r} \cdot \Big(\#\mathcal{S}_{\mathrm{e}}(F_i^q(p), b_1, \dots, b_r) - \#\mathcal{S}_{\mathrm{o}}(F_i^q(p), b_1, \dots, b_r)\Big). \tag{21}$$

where $\delta_{b_1,\dots,b_r}$ is the integer defined by the sequence $b_1, ..., b_r$. Therefore

$$\begin{aligned}
(21) &= \sum_{(b_1,\dots b_r)\in\mathcal{C}} \delta_{b_1,\dots,b_r} \cdot \Big(\#\mathcal{S}_{\mathrm{e}}(F_i^q(p'), b_1, \dots, b_r) - \#\mathcal{S}_{\mathrm{o}}(F_i^q(p'), b_1, \dots, b_r)\Big) \\
&= \sum_{(b_1,\dots b_r)\in\mathcal{C}} \delta_{b_1,\dots,b_r} \cdot \Big(\text{the coefficient of } \kappa_{c_1} \dots \kappa_{c_{q-1}} \text{ in } \varphi \circ \eta(\rho^q(w_i'^{\varepsilon}))\Big) \\
&= \sum_{(b_1,\dots b_r)\in\mathcal{C}} \delta_{b_1,\dots,b_r} \cdot \mu(p'; b_1, \dots, b_r, i).
\end{aligned}$$

Since for any $(b_1, ..., b_r)$ in $\mathcal{C}$, $\mu(p'; b_1, \dots, b_r, i)$ is divided by $\Delta(p'; c_1, ...c_{q-1}, i)$, the above formula is zero modulo $\Delta(p'; c_1, ...c_{q-1}, i)$.

We will show that (20) is equal to (21). Let A_+ and A_- denote $\{G \in \mathcal{K}_+ \mid \hat{G} \in \mathcal{S}(F_i^q(p), b_1, \dots, b_r)\}$ and $\{G \in \mathcal{K}_- \mid \hat{G} \in \mathcal{S}(F_i^q(p), b_1, \dots, b_r)\}$ respectively. Let B_+ and B_- denote $\mathcal{S}_{\mathrm{e}}(F_i^q(p), b_1, \dots, b_r)$ and $\mathcal{S}_{\mathrm{o}}(F_i^q(p), b_1, \dots, b_r)$ respectively. If $A_+ \sqcup A_- = \emptyset$, then we set $\delta_{b_1,\dots,b_r} = 0$. If not, we define a map $\theta : A_+ \sqcup A_- \longrightarrow B_+ \sqcup B_-$ by $\theta(g) = \hat{g}$ for any $g \in A_+ \sqcup A_-$. Then θ is a surjective. In fact there exists h in $B_+ \sqcup B_-$ such that for any g in $A_+ \sqcup A_-$, the image of g under θ is not equal to h, then $A_+ \sqcup A_- = \emptyset$. In addition for any $e_1, e_2 \in B_+$,

$$\begin{aligned}
\#(\theta^{-1}(e_1) \cap A_+) &= \#(\theta^{-1}(e_2) \cap A_+) \\
\#(\theta^{-1}(e_1) \cap A_-) &= \#(\theta^{-1}(e_2) \cap A_-).
\end{aligned}$$

Let us denote β_+ (β_-) the value of the above first (second, respectively) identity. Moreover for any $e_1 \in B_+, e_3 \in B_-$,

$$\begin{aligned}
\#(\theta^{-1}(e_1) \cap A_+) &= \#(\theta^{-1}(e_3) \cap A_-) = \beta_+ \\
\#(\theta^{-1}(e_1) \cap A_-) &= \#(\theta^{-1}(e_3) \cap A_+) = \beta_-.
\end{aligned}$$

Let $\gamma_+ = \#B_+$ and $\gamma_- = \#B_-$.

$$\begin{array}{rcl} A_+ & = & \bigsqcup_{e_1 \in B_+} (\theta^{-1}(e_1) \cap A_+) \bigsqcup_{e_3 \in B_-} (\theta^{-1}(e_3) \cap A_+), \\ A_- & = & \bigsqcup_{e_1 \in B_+} (\theta^{-1}(e_1) \cap A_-) \bigsqcup_{e_3 \in B_-} (\theta^{-1}(e_3) \cap A_-) \end{array}$$

show

$$\begin{array}{rcl} \#A_+ & = & \gamma_+\beta_+ + \gamma_-\beta_-, \\ \#A_- & = & \gamma_+\beta_- + \gamma_-\beta_+. \end{array}$$

Therefore

$$\begin{array}{rcl} \#A_+ - \#A_- & = & (\gamma_+\beta_+ + \gamma_-\beta_-) - (\gamma_+\beta_- + \gamma_-\beta_+) \\ & = & (\beta_+ - \beta_-)(\gamma_+ - \gamma_-) \\ & = & (\beta_+ - \beta_-)(\#B_+ - \#B_-). \end{array}$$

Hence we may set $\delta_{b_1,\ldots,b_r}$ as $\beta_+ - \beta_-$. □

8. Welded links

Fixing α, we define an *H4 move* on nanophrases over α as follows. In the moves on a nanophrases, the lower cases x, y, z and t represent arbitrary sequences of letters, possibly including one or more '|', so that the phrase on each side of the move is nanophrase. The move is

H4-1 move : if $(|A|, |B|) \in T_1$,
$(\mathcal{A}, xAyABzBt) \longleftrightarrow (\mathcal{A}, xAyBAzBt)$.
H4-2 move : if $(|A|, |B|) \in T_2$,
$(\mathcal{A}, xAByAzBt) \longleftrightarrow (\mathcal{A}, xBAyAzBt)$.
H4-3 move : if $(|A|, |B|) \in T_3$,
$(\mathcal{A}, xAyBzABt) \longleftrightarrow (\mathcal{A}, xAyBzBAt)$,

where T_1, T_2 and T_3 are subsets of $\alpha \times \alpha$.

Let T be the triple (T_1, T_2, T_3). We define the *extended M-homotopy* to be the equivalence relation of nanophrases over α generated by isomorphisms, H1,2,3,4 moves with respect to (α, τ, S) and T, self crossing moves with respect to σ and shift moves with respect to ν.

Here we recall that two virtual link diagrams are said to be *welded equivalent* if one may be transformed into the other by a sequence of generalized Reidemeister moves and the upper forbidden moves in Fig. 11. *Welded link* can then be defined to be the welded equivalence class of virtual link diagrams.



Figure 11. Upper forbidden move

Theorem 8.1. *Under the homotopy defined by (α_v, τ_v, S_v), ν_v and T_v, the set of homotopy classes of nanophrases over α_v is in bijective correspondence with the set of ordered welded links, where T_v is the triple (T_1, T_2, T_3) and*

$$T_1 = \{(a_+, a_+), (a_+, b_-), (b_-, a_+), (b_-, b_-)\},$$
$$T_2 = \{(a_-, b_-), (b_+, a_+), (a_-, a_+), (b_+, b_-)\},$$
$$T_3 = \{(a_-, a_-), (a_-, b_+), (b_+, a_-), (b_+, b_+)\}.$$

Proof. By Theorem 2.1, the set of homotopy classes of nanophrases over α_v under the homotopy with respect to (α_v, τ_v, S_v) and ν_v is in bijective correspondence with the set of virtual links. Consider the upper forbidden move. It suffices to consider the case when the orientation of three arcs are as illustrated in Fig. 12. (The deformations involving other orientations of the arcs can be obtained as compositions of this one with isotopy and local deformations of the second Reidemeister moves.) There are 6 cases to consider depending on the order in which one traverses the three arcs involved. Let the order of arcs be illustrated in Fig. 12. Then this move transforms the associated phrase from $xAyBzABt$ to $xAyBzBAt$ where x, y, z and t are words not including the letters A and B, and $|A| = |B| = a_-$. Contrarily if two nanophrases are represented by $xAyBzABt$ and $xAyBzBAt$, the associated virtual link diagrams are related by upper forbidden moves. We can show the others cases in the same way. Therefore under the homotopy defined by (α_v, τ_v, S_v), ν_v and T_v, the set of homotopy classes of nanophrases over α_v is in bijective correspondence with ordered welded links. □



FIGURE 12.

Theorem 8.2. *Let p be an n-component nanophrase. Let $c_1, c_2, \ldots, c_u, i$ be a sequence of integers between 1 and n such that $c_1, c_2, \ldots, c_u, i$ are pairwise distinct. Then $\bar{\mu}(p; c_1, c_2, \ldots, c_u, i)$ is an invariant under extended M-homotopy of nanophrases with respect to (α_v, τ_v, S_v) and T_v associated with the upper forbidden move, ν_v and σ_v.*

Proof. By the proof of Theorem 3.3, it suffices to show that $\mu(p; c_1, \ldots, c_u, i) = \mu(p'; c_1, \ldots, c_u, i)$ if p and p' are related by H4 move corresponding to an upper forbidden move. Suppose $p = xAByAzBt$ and $p' = xBAyAzBt$ where $|A| = |B| = b_-$. (We can show the others cases in the same way.) Since $\eta(A) = \eta(B)$, Theorem 8.2 follows from the same arguments as in proof of Proposition 6.3. □

Note that two nanowords are always equivalent under extended M-homotopy defined by (α_v, τ_v, S_v), ν_v, T_v and σ_v associated with the welded equivalence relation.

ACKNOWLEDGEMENT

The author thanks Professor Sadayoshi Kojima for his valuable suggestions and comments. She want to thank Professor Michael Polyak for informing her his result and giving many valuable comments. Finally, she want to thank Dr. Tomonori Fukunaga for valuable comments.

REFERENCES

[1] H. A. Dye and L. H. Kauffman, *Virtual homotopy*, J. Knot Theory Ramifications **19** (2010), no. 7, 935–960. MR 2673693

[2] T. Fukunaga, *Homotopy classification of nanophrases with less than or equal to four letters*, arXiv:0904.3478. (2009).

[3] A. Gibson, *Factorization of homotopies of nanophrases*, arXiv:0910.5281v2 (2009).

[4] L. H. Kauffman, *Virtual knot theory*, European J. Combin. **20** (1999), no. 7, 663–690. MR 1721925 (2000i:57011)

[5] Olga Kravchenko and Michael Polyak, *Diassociative algebras and Milnor's invariants for tangles*, Lett. Math. Phys. **95** (2011), no. 3, 297–316. MR 2775128

[6] J. Milnor, *Link groups*, Ann. of Math. (2) **59** (1954), 177–195. MR 0071020 (17,70e)

[7] ———, *Isotopy of links. Algebraic geometry and topology*, (1957), 280–306.

[8] V. Turaev, *Curves on surfaces, charts, and words*, Geom. Dedicata **116** (2005), 203–236. MR 2195447 (2007b:57031)

[9] ———, *Knots and words*, Int. Math. Res. Not. (2006), Art. ID 84098, 23. MR 2276346 (2007k:57017)

[10] ———, *Lectures on topology of words*, Jpn. J. Math. **2** (2007), no. 1, 1–39. MR 2295606 (2008i:57024)

[11] ———, *Topology of words*, Proc. Lond. Math. Soc. (3) **95** (2007), no. 2, 360–412. MR 2352565 (2008i:57025)

Department of Mathematics, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan

E-mail address: `kotorii.y.aa@m.titech.ac.jp`